

Abundance, Growth, and Mortality of Selected Oklahoma Blue Catfish Populations: Implications for Management of Trophy Fisheries

Jeff Boxrucker, Oklahoma Fishery Research Laboratory, Oklahoma Department of Wildlife Conservation, 500 E. Constellation, Norman, OK 73072

Kurt Kuklinski, Oklahoma Fishery Research Laboratory, Oklahoma Department of Wildlife Conservation, 500 E. Constellation, Norman, OK 73072

Abstract: Blue catfish *Ictalurus furcatus* electrofishing samples were collected on nine Oklahoma reservoirs and age and growth estimates were made on seven of these reservoirs. Catch rates of blue catfish were high (up to 700/h) on all reservoirs sampled but catch rates of preferred-sized (>762 mm total length) catfish were low (<5/h). Growth rates varied widely both within and among reservoirs but were generally slow with blue catfish not reaching preferred size until ages 13–16. A negative relation between catch rates and growth was identified. Total annual mortality rates averaged 26% for the seven populations sampled. Given growth and mortality rates estimated in this study, only 2%–3% of age-1 blue catfish reach preferred size in Oklahoma reservoirs. A management initiative stressing angler harvest of small individuals while restricting harvest of preferred-sized blue catfish is suggested.

Key words: blue catfish, age and growth, mortality

Proc. Annu. Conf. Southeast. Assoc. Fish and Wildl. Agencies 60:152–156

Angler interest in the pursuit of “trophy-sized” catfish has increased in recent years. Most catfish anglers (71%) take at least one trip annually to pursue trophy catfish (Arterburn et al. 2001). Fishing for trophy-sized fish is more important for blue catfish *Ictalurus furcatus* and flathead catfish *Pylodictis olivaris* anglers than for anglers pursuing channel catfish *Ictalurus punctatus* (Wilde and Ditton 1999, Arterburn et al. 2002). However, the emphasis placed on managing catfish fisheries by agencies appears to be lagging behind angler interest that they do so. Only 2% of agency experts surveyed by Arterburn et al. (2002) indicated that their agencies emphasized managing trophy catfish fisheries even though 75% of catfish anglers surveyed were in favor of developing trophy fisheries. Lack of biological information on catfish populations was the primary reason given for the low emphasis placed on managing catfish fisheries.

Lake Texoma has a reputation as a world-class blue catfish fishery: a former rod and reel world record blue catfish (55.2 kg) was caught there in January 2004. The Oklahoma Department of Wildlife Conservation (ODWC) has been collecting abundance data since the early 1990s but concerns have arisen in recent years that increased fishing pressure on the largest individuals could jeopardize the trophy status of the fishery. Blue catfish growth data were collected on Texoma in 2003 (Mauck and Boxrucker 2005). Growth rates were relatively slow (blue catfish averaged 584 mm at age 10) and highly variable, making management of the fishery for trophy potential challenging. The need to collect growth and mortality information on additional blue catfish populations was

recognized by ODWC staff. The objectives of this study were to: 1) collect abundance, age structure, growth rate, and mortality data on blue catfish populations in selected Oklahoma reservoirs; and 2) use this information to formulate management strategies to enhance and/or preserve the trophy potential of these fisheries.

Methods

Blue catfish abundance (catch rate) data were collected on nine Oklahoma impoundments (four mainstem and five tributary; Table 1). Age and growth samples were collected on seven of these impoundments (catch rate data only were collected on Ft. Cobb and Frederick reservoirs).

Low-frequency (15 pulses/sec), low amperage (4 amps), pulsed-

Table 1. Physical and chemical characteristics of nine Oklahoma reservoirs sampled for blue catfish.

Lake	Year sampled	Surface area (ha)	Mean depth (m)	Secchi (cm)	Conductivity ($\mu\text{S}/\text{cm}$)
Ellsworth ^b	2004	2,258	6	61	475
Eufaula ^a	2005	42,540	8	36	752
Frederick ^b	2005	373	4	15	480
Ft. Cobb ^b	2005	1,653	7	38	470
Hugo ^b	2005	5,343	4	36	90
Kaw ^a	2004	6,871	9	30	450
Keystone ^a	2005	9,520	9	86	3300
Texoma ^a	2003	35,600	11	38	2717
Waurika ^b	2004	4,073	7	64	450

a. Mainstem impoundment

b. Tributary impoundment

DC electrofishing samples, targeting blue catfish, were collected in August 2003–2005. Samples were collected in uplake portions of the reservoir on flats in depths of 3–5 m. The electrofishing boat was manned with a driver and two dippers. In addition, two chase boats, each equipped with a driver and two dippers, assisted in collecting fish. Due to the distance that fish surface from the electrofishing boat, chase boats were essential to maximize collections. The electrofishing boat remained stationary until fish began to surface and then moved slowly in the direction of surfacing fish. Eight units of effort (15 min each) were collected per reservoir for a total of 2 h of sampling effort. Total length (mm) and weight (g) were recorded from all blue catfish collected. Otoliths were removed from a subsample (20 fish/20 mm length group) and processed using methods described by Mauck and Boxrucker (2005). Annular rings were counted by two independent readers. Discrepancies in age determinations were rare but when they did occur, an age was assigned by both readers in concert. Ages were assigned to those fish in the electrofishing samples not aged using reservoir-specific age-length keys (Ketchen 1950).

Data Analysis

Catch per unit effort (CPUE) was calculated as number of blue catfish/15-min units of effort, multiplied by four and expressed as number/h. Catch data were partitioned by total catch and catch of blue catfish ≥ 762 mm (preferred size; Gablehouse 1984) (Table 2). Precision of the CPUE data was expressed as coefficient of variation of the mean ($CV_{\bar{x}}$; Cyr et al. 1992). A $CV_{\bar{x}} = 0.20$ was set as a target level of precision. Pearson correlation was used to compare specific conductance with CPUE ($P \leq 0.05$).

Estimates of mean length at age, von Bertalanffy growth parameters, and total annual mortality (A) using catch curves were derived using the Fisheries Analysis Simulation Tools (FAST)

Table 2. Catch per unit effort (N/h) of all blue catfish in the sample ($CPUE_{total}$) and of preferred-sized blue catfish ($CPUE_{\geq 762}$) with respective precision estimates [coefficient of variation of the mean ($CV_{\bar{x}_{total}}$ and $CV_{\bar{x}_{\geq 762}}$)] and total annual mortality rates (A) of blue catfish from selected Oklahoma reservoirs.

Reservoirs	$CPUE_{total}$	$CV_{\bar{x}_{total}}$	$CPUE_{\geq 762}$	$CV_{\bar{x}_{\geq 762}}$	A
Ellsworth	693.5	0.24	0.5	1.00	0.26
Eufaula	390.0	0.15	0		0.21
Frederick	330.5	0.22	0.5	1.00	
Ft. Cobb	124.0	0.28	3.4	0.54	
Hugo	633.5	0.11	0.5	1.00	0.28
Kaw	294.0	0.15	2.0	0.53	0.30
Keystone	224.0	0.28	3.0	0.42	0.23
Texoma	225.0	0.09	3.0	0.83	0.23
Waurika	490.5	0.09	5.0	0.45	0.32

model (Slipke and Maceina 2000). The von Bertalanffy growth curves for each reservoir were constrained (L_{inf}) by the largest fish in the respective sample (Table 3). Age classes not fully recruited to the gear were eliminated from the catch curve analysis. This varied by population as follows: age 1 eliminated-Eufaula, Hugo, Kaw, Keystone, and Waurika; ages 1 and 2 eliminated-Ellsworth; ages 1, 2, and 3 eliminated-Texoma. Simple linear regression was used to compare mean length at age 10 with CPUE ($P \leq 0.05$).

Results

Electrofishing catch rates ($CPUE_{total}$) ranged from 124.0/h (Ft. Cobb) to 693.5/h (Ellsworth; Table 2). Precision of the $CPUE_{total}$ estimates was generally good with $CV_{\bar{x}_{total}} \leq 0.30$ on all lakes sampled with $CV_{\bar{x}_{total}} \leq 0.20$ (target level) on five of the nine reservoirs sampled (Table 2). Catch rates of preferred-size blue catfish ($CPUE_{\geq 762}$) ranged from 0/h (Eufaula) to 5.0/h (Waurika; Table 2). Precision of the $CPUE_{\geq 762}$ data was poor with $CV_{\bar{x}_{\geq 762}} > 0.40$ on all reservoirs sampled. No relations between specific conductance (Table 1) and total $CPUE_{total}$ (Pearson correlation; $r = -0.49$; $P = 0.18$) nor $CPUE_{\geq 762}$ (Pearson correlation; $r = 0.34$; $P = 0.38$) were found. Annual mortality estimates (A) ranged from 0.21 (Texoma) to 0.32 (Waurika; Table 2) and averaged 0.26.

Growth rates were highly variable, both within and among reservoirs (Table 3). Ranges in length at age often were as much as 50% of the mean (Table 3). Growth rates were poor on Ellsworth, Eufaula, and Hugo with age-10 fish averaging 384.4 mm, 427.3 mm, and 487.0 mm, respectively (Table 3). Growth rates on Kaw, Keystone, Texoma, and Waurika were higher with age-10 blue catfish averaging 610 mm, 636.8 mm, 583.7 mm, and 570.2 mm, respectively (Table 3). However, even on these “faster” growing lakes, it took 13–16 years for blue catfish to average 762 mm (preferred size) and a minimum of 10 years for the fastest growing individual in the sample to reach preferred size (Keystone; Table 3). However, it appears that the growth rates of some individuals in a given population greatly exceed the growth curves (Table 3). The largest fish in the Texoma sample measured 1164 mm, 1087 mm, and 1270 mm and were 13, 16, and 19 years old, respectively. Growth appeared to be density related exhibiting a negative relation between mean length at age 10 and CPUE (Fig. 1; simple linear regression; $r^2 = 0.59$; $P = 0.045$; $N = 7$). von Bertalanffy growth parameters for the seven populations are given in Table 4. All year classes, ages 1–10, were represented in each of the seven reservoirs (Table 3). However, based on the catch curves, younger age classes (1–3) may not have been fully recruited to the sampling gear. Assuming a 26% annual mortality rate and given the growth rates listed in Table 3 (13–16 years to reach preferred size), only 2%–3%

Table 3. Mean length (mm) at age and range of lengths for blue catfish from seven Oklahoma reservoirs.

Age	Ellsworth			Eufaula			Hugo			Kaw			Keystone			Texoma			Waurika		
	N	Length	Range	N	Length	Range	N	Length	Range	N	Length	Range	N	Length	Range	N	Length	Range	N	Length	Range
1	2	165.5	148–183	48	156.0	132–191	721	168.4	92–225	21	174.2	157–194	98	194.5	143–306	30	171.3	138–220	22	184.4	132–198
2	32	185.5	144–205	41	202.8	145–243	212	222.7	138–276	75	232.4	182–255	57	276.5	185–383	21	252.8	133–304	278	194.2	138–277
3	227	222.2	127–269	60	256.4	204–309	167	272.9	172–382	259	272.0	227–365	29	356.2	290–427	23	315.3	250–375	196	262.4	166–369
4	243	235.8	169–288	65	295.1	217–365	20	320.4	298–348	129	324.9	269–438	36	422.0	353–504	42	369.7	319–464	195	292.0	223–389
5	419	253.2	189–311	117	272.5	252–456	45	330.7	254–390	39	400.9	344–502	39	454.6	351–550	32	401.9	335–482	66	321.2	206–488
6	141	273.6	226–322	93	350.5	256–420	44	371	338–418	22	480.7	392–544	62	516.3	440–642	35	439.3	311–555	48	380.8	290–513
7	116	297.9	225–371	71	356.6	259–454	46	411.5	360–498	12	511.2	339–610	24	564.3	467–673	19	459.5	394–529	66	457.6	329–558
8	109	320.9	267–367	39	374.7	296–492	4	449.8	424–492	6	543.7	505–592	26	575.6	487–665	47	496.5	400–645	41	468.4	340–587
9	12	339.2	313–465	32	414.5	340–552	2	474.0	474	9	614.1	572–682	12	611.7	464–660	26	535.9	436–712	35	537.9	424–641
10	22	384.4	341–465	83	427.3	350–566	3	487.0	442–512	1	610		10	636.8	524–857	17	583.7	488–719	10	570.2	383–737
11	57	393.7	355–470	28	472.7	402–579				6	673.2	625–754	5	666.4	596–709	18	572.8	461–829	5	646.4	516–756
12	32	413.5	360–481	34	479.0	384–604				1	737		11	714.4	596–860	12	685.7	477–995	2	750.0	725–775
13	13	382.0	308–487	27	490.6	423–664							1	793		6	881.0	555–1164	4	936.3	864–1050
14	3	554.7	484–670	16	498.1	438–560				5	701.8	633–789				1	696		3	631.3	502–885
15	2	591	456–726	12	486.2	397–622							2	765	717–813	4	851.8	720–994	2	862.0	860–864
16	1	822		7	505.4	440–576				1	830					4	933.0	540–1087			
17				1	531					1	853					1	954				
18	4	600.3	455–783	1	462														1	903	
19							1	926								1	1270				
20				1	504																
21	1	898		2	531.0	496–566															
23	1	896																			
24	1	493																			




Figure 1. Relation of mean length at age 10 (LGTHAGE10) and electrofishing catch rates (CPUE) from blue catfish populations from seven Oklahoma reservoirs. $LGTHAGE10 = 691.56 - 0.3869 * CPUE$ ($r^2 = 0.59$; $P = 0.045$; $N = 7$).

Table 4. von Bertalanffy growth parameters from blue catfish populations from seven Oklahoma reservoirs. L_{inf} = maximum theoretical length (mm) that can be obtained; k = growth coefficient; t_0 = time in years when length would theoretically be equal to 0.

Lake	L_{inf}	k	t_0
Ellsworth	898	0.063	-0.665
Eufaula	622	0.091	-2.526
Hugo	512	0.214	-0.677
Kaw	853	0.136	-0.151
Keystone	860	0.133	-0.919
Texoma	964	0.077	-1.843
Waurika	1050	0.095	0.114

of age-1 blue catfish survive long enough in Oklahoma reservoirs to reach preferred size (Fig. 2).

Discussion

Although the use of low pulse-frequency electrofishing to sample flathead catfish is well documented (Gilliland 1988, Robinson 1994, Cunningham 1995, 2000), published accounts of the use of electrofishing to sample blue catfish are rare. Corcoran (1979) reported that low-frequency pulsed DC current was effective in immobilizing blue catfish and Justus (1996) used electrofishing to collect blue catfish for contaminant monitoring in Mississippi.

Catch rates in this study were high, approaching 700/h in two of the seven reservoirs sampled. Low-frequency electrofishing catch rates in the Rappahannock River, Virginia in 2006 were 4,698/h (Greenlee 2006). The results of this study coupled with the high catch rate reported from Virginia suggest that low-frequency electrofishing is highly effective for collecting large numbers of blue catfish. The precision of the CPUE data (Table 2) was comparable to what the agency obtains for its largemouth bass spring electrofishing data (Oklahoma Department of Wildlife Conservation, unpublished data). As such, we feel that our electrofishing protocol was adequate to compare annual trends in blue catfish abundance. $CPUE_{\geq 762}$ was low, ranging from 0–5 fish/h, suggesting that fish of this size are rare in all populations sampled. However, precision of the $CPUE_{\geq 762}$ data was poor (Table 2) making that data somewhat suspect. Negative bias toward large blue catfish in low-frequency electrofishing samples is also a concern in Virginia (Greenlee 2006) and in Alabama (E.R. Irwin, Auburn University,

personal communication). Some modification of the sampling protocol (e.g., time of year, habitat) may be warranted to decrease the variability in the $CPUE_{\geq 762}$ data and to determine if that sample is representative of the population.

Growth rates reported in this study were moderate relative to those reported in Graham's (1999) review of blue catfish growth rates. However, none of these studies used otoliths to age blue catfish. Given that aging catfish with spines tends to underestimate the age of older individuals (Mayhew 1969, Muncy 1969) direct comparisons between spine-aged and otolith-aged populations may not be valid (Nash and Irwin 1999, Buckmeier et al. 2002). Otolith-based growth rates of the Rappahannock River, Virginia, blue catfish population were similar to those observed in Oklahoma, averaging only 494 mm at age 10 (B. Greenlee, Virginia Department of Game and Inland Fisheries, personal communication). By contrast, otolith-based blue catfish growth estimates from the Santee-Cooper system in South Carolina indicate that age-10 blue catfish average 801 mm (Lamprecht and White 2006). A modified sampling protocol that is more effective in collecting large blue catfish would include an increased number of fast growing individuals in our samples which may increase the estimates of the average length of these older age classes.

The averaged total annual mortality rate estimated in this study (26%) was less than that previously reported for blue catfish populations (Kelley 1969, Hale 1987, Graham and DeiSanti 1999). However, these estimates may be biased. In our study, underestimation of the abundance of the larger fish in the sample would increase the slope of the catch curves and decrease estimates of annual mortality. Underestimation of the age of older fish in studies using spines as an aging method would have the same effect on estimates of annual mortality.

Management Implications

The current blue catfish angler harvest regulation on blue catfish in Oklahoma is a 15-fish daily creel limit, in aggregate with channel catfish, with no length restriction. No commercial harvest of blue catfish is allowed. Given the high total abundance of blue catfish estimated in this study, consistent recruitment, and relatively slow growth, a liberal daily creel limit is warranted. Our evidence of a density dependent growth response also suggests the need for a liberal creel limit.

The low abundance of preferred-sized blue catfish in all reservoirs in this study suggests the need for management efforts aimed at increasing $CPUE_{\geq 762}$. Reducing harvest of preferred-size blue catfish has the potential of increasing abundance of large individuals in the population. Restricting angler harvest to one preferred-sized fish daily, while maintaining or even increasing the existing




Figure 2. Progressive decline in numbers of fish assuming a 26% annual mortality rate and average growth rates of blue catfish in Oklahoma. Oval indicates the ages at which blue catfish reach preferred size.

liberal daily limit of smaller fish should meet the harvest desires of catfish anglers (Wilde and Ditton 1999), reduce the abundance of small fish thereby improving growth rates, and increase the “trophy-potential” of the state’s blue catfish fisheries.

Angler surveys conducted in other states have indicated desires to increase management efforts to preserve/enhance the trophy-potential of blue catfish fisheries (Arterburn et al. 2002, Reitz and Travnichek 2005). ODWC biologists have been contacted by a number of individual anglers expressing similar desires, but a scientific survey of the demographics and desires of catfish anglers in Oklahoma is needed prior to implementing any regulation change aimed at increasing the numbers of large blue catfish in Oklahoma waters.

Acknowledgments

We wish to thank the staffs of ODWC Management Section and the Oklahoma Fishery Research Laboratory for assistance with field collections and otolith removal. We particularly wish to acknowledge Troy Hatfield and Chas Patterson for their expertise in otolith aging. This project was funded through Sport Fish Restoration Grant F-44-D, Project 5.

Literature Cited

- Arterburn, J., D. Kirby, and C. Berry. 2002. A survey of angler attitudes and biologist opinions regarding trophy catfish and their management. *Fisheries* 27(5): 10–21.
- Buckmeier, D.L., E.R. Irwin, R. K. Betsill, and J.A. Prentice. 2002. Validity of otoliths and pectoral spines for estimating ages of channel catfish. *North American Journal of Fisheries Management* 22:934–942.
- Corcoran, M.F. 1979. Electrofishing for catfish: use of low-frequency pulsed direct current. *The Progressive Fish Culturist* 41:200–201.
- Cunningham, K.K. 1995. Comparison of stationary and mobile electrofishing for sampling flathead catfish. *North American Journal of Fisheries Management* 15:515–517. 2000. Influence of environmental variables on flathead catfish electrofishing catch. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 52:125–135.
- Cyr, H., J.A. Downing, S. Lalonde, D.B. Baines, and M.L. Pace. 1992. Sampling larval fish populations: choice of sample number and size. *Transactions of the American Fisheries Society* 121:356–368.
- Gilliland, E. 1988. Telephone, micro-electronic, and generator-powered electrofishing gear for collecting flathead catfish. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 41:221–229.
- Gabelhouse, D.W. Jr. 1984. A length categorization system to assess fish stocks. *North American Journal of Fisheries Management* 4:273–285.
- Graham, K. 1999. A review of the biology and management of blue catfish. Pages 37–49 in E.R. Irwin, W.A. Hubert, C.F. Rabeni, H.L. Schramm Jr., and T. Coon, editors. *Catfish 2000: proceedings of the international ictalurid symposium*. American Fisheries Society, Symposium 24, Bethesda, Maryland.
- and K. DeiSanti. 1999. The population and fishery of blue catfish and channel catfish in the Harry S. Truman Dam tailwater, Missouri. Pages 361–376 in E.R. Irwin, W.A. Hubert, C.F. Rabeni, H.L. Schramm Jr., and T. Coon, editors. *Catfish 2000: proceedings of the international ictalurid symposium*. American Fisheries Society, Symposium 24, Bethesda, Maryland.
- Greenlee, B. 2006. *Catfish management activities: Rappahannock River* *Bulletins* 2006. Virginia Department of Game and Inland Fisheries. Williamsburg.
- Hale, R.S. 1987. Commercial catch analysis, spawning season, and length at maturity of blue catfish in Kentucky Lake, Kentucky-Tennessee. Master’s Thesis. Murray State University, Murray, Kentucky.
- Justus, B. 1996. Observations on electrofishing techniques for three catfish species in Mississippi. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 48:524–532.
- Kelley Jr., J.R. 1969. Growth of blue catfish *Ictalurus furcatus* (LeSueur) in the Tombigbee River of Alabama. *Proceedings of the Annual Conference Southeastern Association of Game and Fish Commissioners* 22:248–255.
- Ketchen, K.S. 1950. Stratified subsampling for determining age distribution. *Transactions of the American Fisheries Society* 79:205–212.
- Lamprecht, S.D. and M.G. White III. 2006. *Fisheries investigations in lakes and streams*. South Carolina Department of Natural Resources. Study Completion Report F63–1–4.
- Mauck, P. and J. Boxrucker. 2005. Abundance, growth, and mortality of the Lake Texoma blue catfish population: implications for management. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 58:57–65.
- Mayhew, J.K. 1969. Age and growth of flathead catfish in the Des Moines River, Iowa. *Transactions of the American Fisheries Society* 98:118–120.
- Muncy, R.J. 1959. Age and growth of channel catfish from the Des Moines River, Boone County, Iowa, 1955 and 1956. *Iowa State Journal of Science* 34:127–137.
- Nash, M.K. and E.R. Irwin. Use of otoliths versus pectoral spines for aging adult flathead catfish. Pages 3091–3316 in E.R. Irwin, W.A. Hubert, C.F. Rabeni, H.L. Schramm Jr., and T. Coon, editors. *Catfish 2000: proceedings of the international ictalurid symposium*. American Fisheries Society, Symposium 24, Bethesda, Maryland.
- Reitz, R.A. and V.H. Travnichek. 2005. An analysis of Missouri catfish angler demographics and opinions for fisheries management alternatives. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 58:66–81.
- Robinson, J.W. 1994. *Sampling procedures to capture flathead catfish and channel catfish*. Missouri Department of Conservation, Fish and Wildlife Research Center, Sport Fish Restoration Project F-1–R-43, Final Report, Columbia.
- Slipke, J.W. and M.J. Maceina. 2000. *Fishery analyses and simulation tools (FAST)*. Auburn University, Auburn, Alabama.
- Wilde, G.R. and R.B. Ditton. 1999. Differences in attitudes and fishing motives among Texas catfish anglers. Pages 395–405 in E.R. Irwin, W.A. Hubert, C.F. Rabeni, H.L. Schramm, Jr., and T. Coon, editors. *Catfish 2000: proceedings of the international ictalurid symposium*. American Fisheries Society, Symposium 24, Bethesda, Maryland.