

## PRELIMINARY STUDIES OF BOBCAT ACTIVITY PATTERNS<sup>a</sup>

J. T. KITCHINGS, Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830

J. D. STORY, Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830

*Abstract.* Home range and activity patterns were determined for one male and one female radio-collared bobcat (*Lynx rufus*) in an eastern Tennessee hardwood forest. Home range of the male was calculated to be approximately 3076 ha while the female utilized 1416 ha. Both bobcats' ranges were larger than previously reported values for the southeast. Measurements of both average net distance traveled per day showed the male moved a statistically significant greater distance than the female. The larger home ranges may be primarily the result of relatively low prey populations in the mountainous terrain of eastern Tennessee as compared to upper coastal plains areas where most of the previous research on southeastern bobcats has been carried out.

Proc. Ann. Conf. S.E. Assoc. Fish & Wildl. Agencies 32: 53-59

In recent years the bobcat has become an economically important fur bearer, in part due to the animal's becoming a substitute for the large spotted cats where fur is no longer available on the international market (Laycock 1978). This change in the bobcat's status has brought an urgent need for more knowledge about the animal's behavior. While progress has been made in gaining this knowledge in some parts of the United States, the bobcat has received relatively little attention in the Southeast in general and virtually no attention in the state of Tennessee.

On the Department of Energy's Oak Ridge Reservation in eastern Tennessee more knowledge of the bobcat's habits was needed for the overall wildlife management program. In March 1978, a study of bobcats of the area was initiated with the radiotagging of an adult female and the subsequent capture and tagging of an adult male. The prime objective of this initial study was to determine the approximate size of home ranges of bobcats on the Oak Ridge Reservation, but we also hoped to obtain information on movement patterns while evaluating the use of radiotracking equipment in the ridge and valley terrain of the Reservation. We gratefully acknowledge the assistance of H. Kitchen at the University of Tennessee School of Veterinary Medicine in Knoxville, in anesthetizing the bobcats in this study.

### MATERIALS AND METHODS

#### Study area.

The Oak Ridge Reservation is located 24.1 km west of Knoxville, Tn. Its southern, western, and eastern boundaries are formed by the Tennessee Valley Authority's (TVA) Melton Hill and Watts Bar Reservoirs on the Clinch River. The city of Oak Ridge forms the northern boundary. The area has been under government control for 30 years and has not been unduly disturbed except for experimental use, regulated forest management, highways, and transmission lines.

The dominant oak-hickory (*Quercus-Carya*) association on the 15,000-ha Reservation contains elements of the mixed mesophytic association commonly found in the adjacent Cumberland Mountains. The oak-hickory association is typified by extensive stands of mixed yellow pine and hardwoods as well as oak and hickory. Yellow poplar (*Liriodendron tulipifera*) often forms nearly pure stands on well-drained bottomlands and lower slopes, while willow (*Salix sp.*) sycamore (*Platanus occidentalis*), and boxelder (*Acer negundo*) border streams and are dominant on poorly drained floodplains. Species

<sup>a</sup>Research sponsored by the Office of Environmental Research, U.S. Department of Energy, under contract W-7405-eng-26 with Union Carbide Corporation. Publication No. 1256, Environmental Sciences Division, ORNL.

more commonly found in the mixed mesophytic association, such as beech (*Fagus grandifolia*), sugar maple, (*A. saccharum*), magnolias (*Magnolia sp.*), buckeye (*Octandra sp.*), and basswood (*Tilia americana*), often occur in coves and sheltered slopes. In addition, approximately 2000 ha of the Reservation were planted in loblolly pine (*Pinus taeda*) between 1947 and 1956 while smaller acreages have since been planted in loblolly pine, black walnut (*Juglans nigra*), river birch (*Betula nigra*), sycamore, and poplar (Kitchings and Mann 1976).

#### Trapping procedures.

Bobcats used in our study were captured in Sullivan live traps using parts of road-killed whitetailed deer (*Odocoileus virginianus*) as bait. Traps were set along gravel backroads where bobcats had been observed. Captured animals were taken to the University of Tennessee School of Veterinary Medicine where they were transferred to a squeeze cage and then anesthetized by an intramuscular injection of Ketamine hydrochloride. The first animal, an adult female, 6.8 kg, caught on 2 March 1978, was equipped with a transmitter and released 4 March 1978, at the capture site. The second bobcat, a 9.5 kg, adult male, caught on 31 March 1978, was measured, ear tagged, equipped with a transmitter, and released at his capture site on 4 April 1978. While the animals were held in captivity, food and water were supplied *ad libitum*.

#### Radio equipment.

Transmitters used in this study were the pulsing, high frequency type in the 150 mHz range, purchased as prepared units from Wildlife Materials Incorporated. Battery life for these units is listed by the manufacturer as 555 days + 111 days.

A single receiver was used with a truck-mounted whip antenna and a hand-held 4-element Yagi antenna. Range of reception varied and while signals were received at distances up to 3.8 km on both the whip and Yagi antennas, most bearings were taken at distances of 1.6 km or less from the transmitters. All tracking was done from the ground, and there were a few days when we were unable to locate the bobcats. On the few occasions when we were able to obtain only 1 bearing for an animal, no attempt was made to estimate the distance to the transmitter along this 1 bearing. We found that signal strength varied tremendously depending on the cover afforded the bobcat by its immediate surroundings.

#### Radiotracking procedure.

The general location of a bobcat was determined using the truck-mounted whip antenna and driving until the signal from 1 of the bobcats was received. This antenna also enabled us to select sites of optimum signal strength before trying to determine the magnetic bearing to the transmitter. Once a site was selected, the truck-mounted antenna was disconnected from the receiver and the hand-held Yagi was connected. The direction of greatest signal strength was then determined using the Yagi antenna, and the magnetic bearing was read from a Silva Type 1 hand-held compass.

Magnetic bearings were plotted on a 1:24,000 scale map using a Jeppesen Navigation Plotter. While 2 such plots separated by a sufficiently wide angle (preferably 45<sup>m</sup>–135<sup>m</sup>) showed the approximate location of the bobcat, frequently a third bearing was taken to verify the accuracy of the first 2. Practice sessions with the 2 transmitters used in this study and with other similar transmitters demonstrated that our radio-locations were 148 + 30 m (N = 23) within actual location.

#### Data Analysis

Home range areas were determined by the convex polygon method to facilitate comparisons of our data with those of other investigators. Jennerich and Turner (1969) pointed out that the convex polygon method severely underestimated the home range, particularly when the sample size was low. However, based on subsequent calculations by




Fig. 1. Home range estimates for a male and a female bobcat radio-tracked on Oak Ridge Reservation from March through May 1978.

Gardner (1978) the sample sizes in our study and those in the literature are large enough that the convex polygon estimation is almost the same as the area estimation achieved by use of Jennerich and Turner's approach. To compare daily movements between the male and female, the linear distance between 5 or more consecutive daily locations was measured. The "student's" *t* test for was used to determine statistical significance between the mean of the 2 samples.

## RESULTS

### Home range.

The total area covered by each of the bobcats during the 4 March–23 May tracking is shown in Fig. 1. The female was located 159 times during 68 tracking-days from 4 March 1978 through 23 May 1978, while we located the male 83 times during 34 tracking-days from 4 April 1978 through 17 May 1978. Area used by the female and male were calculated to be 1416 ha and 3076 ha respectively. The ratio of the area covered by the male to that used by the female was 2.2 and overlaps 90% of the female's range (Fig. 1). Visual observations of a third animal near the northeast perimeter of the male's territory indicate that at least 1 other animal is in close proximity to the 2 currently being radiotracked, but attempts to trap this individual have been unsuccessful.

## Movements.

The tracking data revealed that the animals were generally most active during the time between 1700 and 1100 hours. The female utilized easily recognizable daytime resting areas up until the time she chose a permanent den site. After recording daily positions for the female during the first 2 weeks after capture, we spent two 5 day periods 20-24 and 27-31 March taking periodic measurements of the animal's position from 0400 to 2400 and 1700 to 0100 hours. The number of locations per day varied from 3 to 13. During the first 5 days (20-24 March) the female appeared to spend most of her time in the area outlined in Fig. 2, Map A. The daytime resting areas were consistently used during this period. Tracking during the second 5 days indicated that the female had moved eastward (Fig. 2, Map B) and established a permanent den site. Previous daytime rest areas apparently were abandoned once the major den site was determined. Subsequent daily tracking during the month of April confirmed the continued use of the area as a den site.




Fig. 2. Areas used by a female bobcat radio-tracked on Oak Ridge Reservation during three five-day periods of intensive radiolocation efforts.

Map A: Area used March 20 through March 24

Map B: Area used March 27 through March 31

Map C: Area used May 1 through May 5

In the first week of May, continuous tracking was again undertaken (Fig. 2, Map C). The area covered by the female bobcat during the 27-31 March and 1-5 May periods was 259 ha and 235 ha, respectively, approximately 18% and 17% of the total home range measured for the entire period from March to May. The existence of a number of temporary dens was indicated. On 2 occasions the female was disturbed by our presence in the area near the major den site.

Data accumulated on the male bobcat since his release do not indicate the use of a number of well-established daytime resting areas. Nor did the male seem to restrict his movements to any particular area during the period 1-5 May when continuous measurements of his movements were made in conjunction with those of the female. In fact, he was located virtually throughout his range (Fig. 1) during the tracking effort.

While the number of locations per day varied from 3 to 13, net distance between daily locations always was calculated from the first location of each (whether the cat was located 3 or 13 times). Measurements of net distance between daily locations showed the male moving a significantly ( $P < 0.001$ ;  $t = 126.8$ , d.f. 84) greater distance, averaging a net distance of 3.3 km per day, while the female covered 1.0 km daily. The actual distance that the cats moved on a daily basis is much greater than indicated by daily average net distance. Determinations of average net distance moved by a cat during a day were based only on days on which we obtained 5 or more locations; the female measurements were based on 96 locations on 13 days, while the male measurements used 38 locations on 4 days since we could not actually measure the circuitous routes taken by the moving cats. On days when 5 or more consecutive locations were made, the female moved an average distance of 4.0 km (range 1.2 - 8.5 km) while the male covered an average distance of 8.8 km (range 5.8 - 11.7 km). These values are significantly different at  $P < 0.001$  ( $t = 6.41$ , d.f. = 14).

Even though no direct observations of hunting by either bobcat were made, the location of animals during the active period pointed to extensive use of early successional vegetation. Maintained experimental old-fields, power-line rights-of-way and roadsides comprised the majority of locations, with swamp-like floodplain and cutover areas around nuclear waste disposal areas also being used for hunting. These types of vegetational communities would, of course, be prime areas for both rabbits (*Sylvilagus floridanus*) and cotton rats (*Sigmodon hispidus*).

It was not unusual to find either animal moving during the daytime. However, most hunting is believed to have been done at night. Vegetative cover was so dense in the areas used by the bobcats that neither animal was observed during radio-tracking operations.

Adult interactions.

We found the male and female together in the vicinity of the den site on 7 different occasions. At no time during our night tracking did we find any indication that the male and female worked together during hunting efforts.

## DISCUSSION

The area used by the 2 animals reported in our study are considerably larger than the home ranges calculated for the southeast area by Marshall and Jenkins (1966) and by Miller (personal communication, Auburn Univ., 1978). The former calculated ranges of 247 and 463 ha for 2 females and 347 ha for a young male, while Miller reported females utilizing 202 ha and males twice that — 405 ha. In both cases the study sites were composed of larger areas of old-field type vegetation, areas where prey species would be abundant, thus limiting the necessity for extensive excursions for food. The large range of our 2 cats probably reflects the low food base available in the area. Of the total acreage included in the range calculation for both sexes, there is less than 40.5 ha of old-field. While we have no data for prey abundance since these studies are just getting under way, informal discussions with a colleague working with cotton rats indicated that population values are

low possibly as a result of the severe 1977-78 winter. Likewise, visual observations of cottontail rabbits along roadsides do not indicate an abundance of this species either. We feel that the usage area of the bobcats is indicative of those areas in the Ridge and Valley Province where vegetation is principally forest with only a small proportion in active or abandoned farmland.

While small game, such as rabbits and rodents, comprise a large portion of the bobcat's foodstuff, they are by no means the only food source. Literature on bobcat food habits indicate that remains of whitetail deer are a commonly found component of the cat's diet (Marston 1942, Rollings 1945, Progulske 1955, Pollack 1951, and Young 1958). A report on the bobcat in Tennessee (Buttrey 1978) indicated that 20% of stomach content samples taken from animals in the Cumberland Plateau area contained remains of whitetail deer; while more deer was found in bobcat stomachs during hunting season than during spring and summer, and most deer probably was eaten as carrion, Buttrey (1978) felt that "regardless of how acquired, deer constitutes a substantial portion of the bobcat's diet." Reports of bobcats attacking and killing adult deer are common (Marston 1942, and McCord 1974) and it is possible that this species provides food for the bobcats on the Oak Ridge area. Deer are abundant on the Reservation since hunting is prohibited and are going through a stage of rapid population growth thus making this species more available to the cats as a food source. The interaction between deer and bobcat warrants further investigation.

Bailey (1974) also found the male bobcat's range to be 2.2 times greater than that of the female. As he pointed out, survival of the offspring is highly dependent on the female's successful rearing of the kittens and thus she must be less mobile than the male. This is supported by our finding a significant difference in average net distance moved daily by males and females and in linear distance between day-to-day locations.

Various studies (Marshall and Jenkins 1966, Bailey 1974) have noted a close proximity between sexes on occasion but no long term interaction. None of the literature we surveyed indicated that males have any rearing responsibilities and none of our data supports any conclusion to the contrary.

We believe that in a situation like that at Oak Ridge, there probably exists a firm territorial system with a set spatial arrangement of resident bobcats as described by Marshall and Jenkins (1966) and Bailey (1974); the protection from hunting probably affords resident bobcats a considerable degree of permanence regarding their social organization (home range, territory, interactions with other bobcats) not enjoyed by animals in hunted areas. Consequently it may very well be that the Reservation acts as a nursery or reservoir for this species. As young reach an age of independence and leave the mother they may move off the Reservation into areas where they are then accessible to trappers and hunters just as Bailey found young leaving the protected area in his Idaho study and then being captured by fur trappers.

#### LITERATURE CITED

- Bailey, T. N. 1974. Social organization in a bobcat population. *J. Wildl. Manage.* 38(3):435-446.
- Buttrey, G. 1978. The bobcat's niche in nature. *Tenn. Wildl.* 1(4):17-18.
- Jennerich, R. I., and F. B. Turner. 1969. Measurement of non-circular home range. *J. theoret. Biol.* 22:227-237.
- Kitchings, T., and L. K. Mann. 1976. A description of the terrestrial ecology of the Oak Ridge Environmental Research Park. ORNL/TM-5073. Oak Ridge National Laboratory, Oak Ridge, Tennessee. 58 pp.
- Laycock, G. 1978. Bobcat embargo is short-lived. *Audubon* 80(2):108.

- Marshall, A. D., and J. H. Jenkins. 1966. Movements and home ranges of bobcats as determined by radiotracking in the upper coastal plain of west-central South Carolina. Proc. Southeastern Assoc. Game and Fish Commissioners 20:206-214.
- Marston, M. A. 1942. Winter relations of bobcats to white-tailed deer in Maine. J. Wildl. Manage. 6(4):328-337.
- McCord, C. M. 1974. Selection of winter habitat by bobcats (*Lynx rufus*) on the Quabbin Reservation, Massachusetts. J. Mammal. 55(2):428-437.
- Pollack, E. M. 1951. Food habits of the bobcat in the New England states. J. Wildl. Manage. 15(2):209-213.
- Progulske, D. R. 1955. Game animals utilized as food by the bobcat in the Southern Appalachians. J. Wildl. Manage. 19(2):249-253.
- Rollings, C. T. 1945. Habits, foods and parasites of the bobcat in Minnesota. J. Wildl. Manage. 9(2):131-145.
- Young, S. P. 1958. The bobcat of North America. Stackpole Co., Harrisburg, Pennsylvania. 193 pp.